

COMMENTS ON THE ARTICLE: “DEPRESSION OF THE PERUVIAN ELDERLY ADULT AND ASSOCIATED SOCIODEMOGRAPHICAL VARIABLES, ANALYSIS OF ENDES 2017”

COMENTARIOS SOBRE EL ARTÍCULO:
“DEPRESIÓN DEL ADULTO MAYOR PERUANO Y VARIABLES SOCIODEMOGRÁFICAS ASOCIADAS:
ANÁLISIS DE ENDES 2017”

Jaime Lama-Valdivia^{1,2,a,b}, Rodrigo Rodríguez-Del Pozo^{2,a}

Mr. Editor

In the article published by Baldeón-Martínez et al⁽¹⁾ on depression in the elderly, the authors estimated a prevalence of depression of 14.2% in a probabilistic sample of 4917 people over 60 years of age in our country. Factors associated with this diagnosis were female sex, lack of education, age over 75, living in rural areas and being poor. This study was carried out on the basis of data from the 2017 Demographic and Family Health Survey (DHS), using the instrument Patient Health Questionnaire (PHQ-9) for the diagnosis of depression. The study presents some strengths such as the broad coverage, the quality of sampling and the detailed and up-to-date information on respondents, especially on socio-demographic aspects.

However, its analysis allows us to find a series of limitations and biases that the authors do not comment on in the discussion, such as: the methodology and objective of the study, the validity of the instrument used in the geriatric population and the relevance or redundancy of the research in relation to previously published works on the topic.

The specific objective of the DHS was not to estimate the prevalence rate of depression in the Peruvian elderly, being the information of the article obtained from a secondary database. On the other hand, the authors mention as an objective to determine the variables associated with geriatric depression, using the formula of sample size of frequencies used by the DHS and not one of the risk factors, which would be most appropriate and would allow a correct interpretation of the Odds Ratio (OR) obtained.

The instrument used was the PHQ-9 validated in Peru by a committee of mental health experts, but as Calderón et al⁽²⁾ mentioned in their 2012 article, validation had not been done in the population, where there could be socio-cultural differences between coastal, highlands and jungle communities. As the instrument supplemented by self-report, the authors do not mention the criteria for exclusion from their study, such as chronic cognitive impairment, delirium, hearing loss, decreased visual acuity, locomotor system disorder or language barrier, all common conditions in elderly⁽³⁾.

Recent revisions on the instruments used in the diagnosis of depression in the elderly, agree that the Geriatric Depression Scale (GDS) or Yesavage Scale and the Center for Epidemiological Studies-Depression Scale (CES-D) are the most reliable instruments for their use, taking into account specific aspects of the presentation of this pathology in this age group, such as attention deficit and memory deficit, somatizations, anxiety and sleep disorder⁽⁴⁾. In fact, in the discussion of their work, the authors compare their findings with those obtained in studies using these instruments.

¹ Faculty of Human Medicine "Manuel Huamán Guerrero", Ricardo Palma University, Lima-Peru.

² National Hospital Hipólito Unanue, Lima-Peru.

^a Geriatrician doctor.

^b Associate professor.

Cite as: Jaime Lama-Valdivia, Rodrigo Rodríguez-Del Pozo. Comments on the article: “Depression of the peruvian elderly adult and associated sociodemographical variables, analysis of endes 2017”. Rev. Fac. Med. Hum. January 2020; 20(1):166-167. DOI 10.25176/RFMH.v20i1.2560

Journal home page: <http://revistas.urp.edu.pe/index.php/RFMH>

Article published by the Journal of the Faculty of Human Medicine of the Ricardo Palma University. It is an open access article, distributed under the terms of the Creative Commons License: Creative Commons Attribution 4.0 International, CC BY 4.0 (<https://creativecommons.org/licenses/by/4.0/>), which allows non-commercial use, distribution and reproduction in any medium, provided that the original work is duly cited. For commercial use, please contact magazine.medicina@urp.pe

Finally, the study does not show greater relevance and is even redundant compared to what was published by Martina et al⁽⁵⁾, who examine depression and associated factors in the Peruvian elderly population according to DHS 2014-2015, finding a prevalence of depression of 14.3% and association with factors such as the female sex, being over 75 years old, lack of education, living without a partner, living in rural areas, poverty and physical disability.

We believe that there is a need for more population studies on the subject, addressing depression in the elderly as a disease, with a focus not only on social and demographic, but also on medicine. One proposal could be to study the groups identified as high-risk, along with a comprehensive geriatric assessment, which would allow for timely intervention spaces⁽⁶⁾.

Authorship Contributions: The authors participated in the generation, collection of information, writing and final version of the original article.

Financing: Self-financed

Interest conflict: The authors declare no conflicts of interest in the publication of this article.

Received: November 15, 2019

Approved: December 26, 2019

Correspondence: Jaime Lama-Valdivia.

Address: Av. Alfredo Benavides 5440, Santiago de Surco, Lima-Perú.

Telephone: 999093383

E-mail: jlamav@hotmail.com

BIBLIOGRAPHIC REFERENCES

1. Paola Baldeón-Martínez, Consuelo Luna-Muñoz, Sandra Mendoza-Cernaqué, Lucy E. Correa-López. Depresión del adulto mayor peruano y variables sociodemográficas asociadas: análisis de endes 2017. Rev. Fac. Med. Hum. Octubre 2019; 19(4):47-52. DOI 10.25176/RFMH.v19i4.2339
2. Calderón M, Gálvez-Buccollini JA, Cueva G, Ordoñez C, Bromley C, Fiestas F. Validación de la versión peruana del PHQ-9 para el diagnóstico de depresión. Rev Peru Med Exp Salud Publica [Internet]. diciembre de 2012 [citado el 1 de septiembrede2019];29(4):5789.Disponibleen:http://www.scielo.org/scielo.php?script=sci_arttext&pid=S172646342012000400027&lng=es&nrm=iso&tng=es
3. Varela L, Chávez H, Herrera A, Méndez F, Gálvez M. Perfil del adulto mayor INTRA II. Lima: Ministerio de Salud; 2004. 133p.
4. Watson LC, Pignone MP. Screening accuracy for late-life depression in primary care: a systematic review. J Fam Pract. 2003; 52(12): 956-64.
5. Martina M, Ara MA, Gutiérrez C, Nolberto V, Piscoya J. Depresión y factores asociados en la población peruana adulta mayor según la ENDES 2014-2015. An la Fac Med [Internet]. el 23 de marzo de 2018 [citado el 1 de septiembre de 2019];78(4):393.Disponibleen:http://revistasinvestigacion.unmsm.edu.pe/index.php/anales/article/view/14259
6. Gálvez-Cano M, Chávez-Jimeno H, Aliaga-Díaz E. Utilidad de la valoración geriátrica integral en la evaluación de la salud del adulto mayor. Rev Peru Med Exp Salud Publica. 2016;33(2):321-7.

Indexed in:



<https://alicia.concytec.gob.pe/vufind/>

