

Biliary ileus: case report

Íleo biliar: caso clínico

Fernanda Kreve¹, Jonas Takada², Janaina Gatto¹, Francisco S. Loss¹, Everson L. A. Artifon²

¹ Medical School Academic, Centro Universitário Fundação Assis Gurgacz. Cascavel, Brazil.

² Department of Surgery, University of Sao Paulo. Sao Paulo, Brazil.

Recibido: 14-8-2016

Aprobado: 2-10-2016

ABSTRACT

Biliary ileus is a uncommon cause of mechanical bowel obstruction, affecting older adult patients who often have other significant medical conditions. It is caused by intestinal impaction of a gallstone that enters the bowel via a cholecysto-enteric fistula. The mortality rate is considerable, ranging between 12 and 27%. Treatment in most cases is surgical, aimed at the resolution of the intestinal obstruction. We report the case of a 55 year old patient diagnosed with biliary ileus, which evolved without complications after a enterolithotomy. We have concluded, based on the literature that an early diagnosis associated with appropriate therapy can lead to a better prognosis.

Keywords: *Ileus; Gallbladder; Intestinal obstruction (source: MeSH NLM).*

RESUMEN

El íleo biliar es una causa poco frecuente de obstrucción intestinal mecánica, que afecta pacientes de edad avanzada que generalmente tienen comorbilidades significativas. Es causada por la impactación de un cálculo biliar en el intestino, que entra en el tracto digestivo por medio de una fístula colecisto-entérica. La tasa de mortalidad es considerable, que va de 12 a 27%. El tratamiento en la mayoría de los casos es quirúrgico y destinado a la resolución de la obstrucción intestinal. Presentamos el caso de una paciente de 55 años diagnosticada con íleo biliar, que evolucionó sin complicaciones después de enterolitotomía. Llegamos a la conclusión, con base en la literatura, que un diagnóstico precoz asociado con el tratamiento adecuado conduce a un mejor pronóstico.

Palabras clave: *Ileus; Vesícula biliar; Obstrucción intestinal (fuente: DeCS BIREME).*

INTRODUCTION

Cholelithiasis is one of the most common surgical diseases. Its incidence is age - related, with increasing prevalence in the elderly population. In its evolution, approximately 2% of patients develop symptoms or complications, such as acute cholecystitis, pancreatitis and choledocholithiasis. In rare cases a cholecysto-enteric fistula may occur, which can lead to intestinal obstruction after migration and impaction of gallstones, creating a condition known as biliary ileus. We report a case of 55 year old female who presented with small bowel obstruction owing to large gallstone in ileum.

CASE REPORT

Patient female, 55 years. Previous history of hypertension and total hysterectomy for benign disease. Reported intense pain in the right hypochondrium for 6

days associated with vomit. On the 3rd day of evolution, the pain started to have diffuse pattern, associated with feculent vomit, abdominal distension and stop elimination of gas and faeces.

On physical examination the patient was hemodynamically stable, anicteric, with mild signs of dehydration, the abdomen was distended with very active bowel sounds and painful diffusely. Leukogram pointed slight elevation, liver and canalicular enzymes (total bilirubin, direct and indirect bilirubin, alkaline phosphatase, gamma glutamyl transferase, transaminase glutamic oxaloacetic and transaminase glutamic pyruvic) within normal parameters.

Computed tomography indicated aerobilia, important distension of the small bowel, calcified and impacted foreign body in the distal ileum, suggesting biliary ileus (Figures 1 and 2). Given the findings, the patient underwent upper digestive endoscopy, which



Figure 1. Computed tomography (axial section) indicating the presence of aerobilia.

identified the existence of wide cholecysto-duodenal fistula, allowing internal visualization of gallbladder, discarding the presence of remaining gallstones (Figures 3 and 4).

Based on the clinical and laboratory tests, the patient was taken to the operating room. Intraoperatively, was observed significant distension of the small bowel up to the region of the distal ileum, which had an intraluminal foreign body, about 60 cm from the ileocecal valve. The gallbladder was partially viewed, and was well adhered to the first duodenal portion, forming a single block inflammatory which did not compromise the structures of the hepatic hilum. The enterotomy was performed on the antimesenteric border of the obstruction, with subsequent extraction of the 5 cm gallstone and enteroraphy (Figures 5 and 6). We chose not to address the gallbladder. Postoperatively, the patient recovered uneventfully and was discharged from the hospital on the fourth day for outpatient treatment.

Exploratory laparotomy was done. Intra-operatively, a gallstone of size 5x3 cm was found obstructing the proximal ileal lumen, causing dilatation of proximal bowel loops and distal collapse. The gallbladder was



Figure 3. Endoscopic view of the duodenal bulb indicating the presence of cholecysto-duodenal fistula.



Figure 2. Computed tomography (sagittal section) showing a foreign body impacted in the ileum.

partially viewed, and was well adhered to the first duodenal portion, forming a single block inflammatory which did not compromise the structures of the hepatic hilum. Proximal longitudinal enterolithotomy was done, with subsequent decompression after gallstone extraction (Figures 5 and 6). We chose not to perform the cholecystectomy and the fistula repair. Postoperatively the patient recovered uneventfully and was discharged on the fourth day for outpatient treatment.

DISCUSSION

Biliary ileus was first described by Bartholin in 1645 during an autopsy ⁽¹⁾. It is a rare condition, being unusual before 50 years and more frequent in women, with an average rate of 3.5: 1 ⁽²⁾. Happens in 1 to 3% of small bowel obstruction in patients up to 65 years of age, incidence which rises to 25% in patients over this age ⁽³⁾. It is a disease presents in only 0.06% of patients with cholelithiasis. It results from repeated



Figure 4. Internal endoscopic view of gallbladder discarding the presence of remaining gallstones.



Figure 5. Gallstone extraction through enterotomy in antimesenteric.

inflammatory conditions in the gallbladder, leading to the involvement of surrounding structures, formation of fistulous orifices and further migration of gallstones to the gastrointestinal tract⁽⁴⁾. The literature describes three main possible fistulas between the biliary tract and digestive tract: cholecysto-duodenal (65-77%), cholecysto-colics (10-25%) and cholecysto-gastrics (5%)⁽⁵⁾. Consequently, the symptoms become varied, as the gallstone may lead to obstruction of any segment of the digestive tract, proximal and distal.

Interestingly, one third of the patients are absent of prior symptoms of biliary colic and up to 50% have no history of gallbladder disease. The occurrence of severe pain located mainly in the right upper quadrant may be the harbinger of the biliary-digestive fistula⁽⁶⁾. After the formation of the fistula and gallstones migration, many symptoms tend to alleviate until complete disappearance. If the gallstone is small, can be eliminated by faeces, leading to a lack of knowledge of the disease⁽⁷⁾. However, relief of symptoms can be only momentary (a few hours or a few days), with subsequent migration of the pain due to the "tumbling phenomenon": migration and impaction of the gallstone into a most distal segment of the digestive tract⁽⁸⁾.

The distal ileum, due to the small size and weak peristalsis, is the most frequent site of obstruction⁽⁹⁾. The symptoms in this case are typical of small bowel obstruction, including nausea, vomiting, distension, dehydration and abdominal pain. The Riegler's triad (aerobilia, distended loops and ectopic gallstone), set of highly suggestive biliary ileus signals, is observed in only 15% of the radiographs, which leads many patients to surgery without a predetermined diagnosis. Aiming to a more accurate diagnosis, computed tomography (CT) can demonstrate Riegler's triad up to 80% of cases⁽¹⁰⁾. Therefore, the image provided by CT is crucial not only for the diagnosis of biliary ileus, but also to helping in the therapeutic decision⁽¹¹⁾.



Figure 6. Gallstone measuring 3 cm x 5 cm.

Another rare complication of cholelithiasis is Mirizzi's syndrome, characterized by extrinsic compression of the bile duct by the infundibulum of the gallstone or cystic duct⁽¹²⁾, which may culminate in cholecysto-biliary fistula. By involving different stages of the same disease, when identified cholecysto-enteric fistula, the possibility of association with Mirizzi's syndrome should be considered. In a study⁽¹³⁾, where patients underwent to a cholecystectomy in a 11 - year period, 29% had cholecysto-enteric fistula associated with Mirizzi's syndrome.

Extracorporeal lithotripsy shock waves for fragmenting the stones, mechanical dilation of stenotic segments and endoscopic removal, have been described as alternatives forms of non-surgical treatment⁽¹⁴⁾. But, these treatments take into account the location of the obstruction, according as the gallstone can be removed by endoscopy or colonoscopy if they are in a favorable position⁽¹⁵⁾. Furthermore, the availability and cost of materials have to be considered too.

The treatment is most often surgical and can be divided into two steps: one is mandatory and for the relief of intestinal obstruction (enterolithotomy); and the other, that is not always possible and often not recommended, is for the cholecystectomy and fistula closure. A more complex and slow operation may result in higher complications rates. Thus, cholecystectomy and fistula repair should be reserved only for selected patients with absolute indications, including acute cholecystitis, gangrene of the gallbladder or residual gallstones. Furthermore, recurrent episodes of biliary ileus are rare, reported in only 5% of patients, and attributed to the residual gallstones previously undetected. When there are no residual gallstones, most fistulas will close spontaneously⁽¹⁶⁾. Therefore, considering the rare recurrence of biliary ileus and excellent tolerability of biliodigestive fistula that does not close spontaneously, just enterolithotomy proves advantageous for most patients with biliary ileus⁽¹⁷⁾.

Being an uncommon cause of small bowel obstruction, the diagnosis of this pathology is usually delayed, sometimes being performed only at the surgery⁽⁷⁾. The late diagnosis worsens the prognosis, with a mortality ranging from 12 to 27%. However, the worst outcome is mainly due to advanced age and their associated comorbidities, which increase the surgical risk. Therefore, it is extremely important to choose the most advantageous and the safe surgical approach. A longer operation time associated with an additional surgical trauma caused by the fistula repair cannot be beneficial for the majority of these patients⁽¹⁸⁾.

In this case, the treatment showed excellent results without associated complications. The fistula closure was not done, following the guidelines described in the literature that demonstrate high rate of complications and small benefit in fistula repair. Moreover, endoscopic visualization of the gallbladder, discarding the presence of residual gallstones, eliminated the possibility of recurrence of biliary ileus in this patient.

Conflicts of Interest: None.

BIBLIOGRAPHIC REFERENCES

- Masannat Y, Shatnawei, A. Gallstone ileus. A review. *Mt Sinai J Med.* 2006;73(8):1132-4.
- Nakao A, Okamoto Y, Sunami M, Fujita T, Tsuji T. The oldest patient with gallstone ileus: report of a case and review of 176 cases in Japan. *Kurume Med J.* 2008;55(1-2):29-33.
- García OS, González CHQ, Téllez-Ávila A. Fístula bilioentérica com impactación de lito gigante em yeyuno. *Rev Gastroenterol Mex.* 2008;73(4):235-8.
- Szajn bok I, Lorenzi F, Rodrigues Júnior AJ, Zantut LF, Poggetti RS, Steinman E, et al. Gallstone ileus resulting in strong intestinal obstruction. *Sao Paulo Med J.* 1995;1(113):721-5.
- Campelo MRO, Menegola VM, Galdino JPC. Íleo biliar: um relato de caso. *Revista AMRIGS* 2015;59(1):35-8.
- Berger MY, Van DerVelden JJ, Lijmer JC, DeKort H, Prins A, Bohnen AM. Abdominal symptoms: do they predict gallstones? A systematic review. *Scand J Gastroenterol.* 2000;35(1):70-6.
- Michele D, Luciano G, Massimiliano F, Stefano R, Roberta D, Ernesto S, et al. Usefulness of CT-scan in the diagnosis and therapeutic approach of gallstone ileus: report of two surgically treated cases. *BMC Surg.* 2013;13 Suppl 2:S6. 8. Reisner RM, Cohen JR. Gallstone ileus: a review of 1001 reported cases. *Am Surg.* 1994;60(6):441-6.
- Echenique Elizondo M, Amondaraín Arratibel JA, Lirón de Robles Sanz C. Íleo Biliar. *Ver Esp Enferm Dig.* 2007;99(11):672-80. 10. Roothans D, Anguille S. Rigler triad in gallstone ileus. *CMAJ.* 2013;185(14):E690.
- Yu CY, Lin CC, Shyu RY, Hsieh CB, Wu HS, Tyan YS, et al. Value of CT in the diagnosis and management of gallstone ileus. *World J Gastroenterol.* 2005;11(14):2142-7.
- Gómez MA, Meneses JC. Usefulness of endoscopy for diagnosing Mirizzi syndrome. *Rev Col Gastroenterol.* 2010;3(25):312-5.
- Beltran MA, Csendes A, Cruces KS. The relationship of Mirizzi syndrome and cholecystoenteric fistula: validation of a modified classification. *World J Surg.* 2008;32(2):2237-43.
- Pezzoli A, Maimone A, Fusetti N, Pizzo E. Gallstone ileus treated with non-surgical conservative methods: a case report. *J Med Case Rep.* 2015;9:15.
- Chatterjee S, Tamonas C, Goutan G, Ambar G. Gallstone ileus an atypical presentation and unusual location. *Int J Surg.* 2008;6(6):55-6.
- Vagefi PA, Ferguson CM, Hall JF. Recurrent gallstone ileus: third time is the charm. *Arch Surg.* 2008;143(11):1118-20.
- Rodríguez-Sanjuán, JC, Casado F, Fernández MJ, Morales DJ, Naranjo A. Cholecystectomy and fistula closure versus enterolithotomy alone in gallstone ileus. *Br J Surg.* 1997;84(5):634-7.
- Doko M, Zovak M, Kopljár M, Glavan E, Ljubicic N, Hochstädter H. Comparison of surgical treatments of gallstone ileus: preliminary report. *World J Surg.* 2003;27(4):400-4.

Correspondence:

Everson L. A. Artifon

Rua Guimaraes Passos, 260/apto 121 Sao Paulo. ZC: 04107-030; Brazil
E-mail: eartifon@hotmail.com