

MATERIAL SUPLEMENTARIO

Anexo 1. Proporciones de resistencia basal, a los tres y seis meses en dos comunidades rurales de Moyobamba y Urubamba.

Antibiótico	Moyobamba			Urubamba		
	Basal (n=47) n (%)	3 meses (n=32) n (%)	6 meses (n=20) n (%)	Basal (n=39) n (%)	3 meses (n=27) n (%)	6 meses (n=14) n (%)
Cotrimoxazol	18 (38,3)	16 (50,0)	10 (50,0)	21 (53,8)	15 (55,6)	8 (57,1)
Ampicilina	22 (46,8)	15 (46,9)	8 (40,0)	21 (53,8)	14 (51,9)	4 (28,6)
Ácido nalidíxico	11 (23,4)	8 (25,0)	10 (50,0)	14 (35,9)	11 (34,4)	5 (35,7)
Azitromicina	6 (12,8)	7 (21,9)	4 (20,0)	5 (12,8)	5 (18,5)	1 (7,1)
Ciprofloxacina	3 (6,4)	4 (12,5)	1 (5,0)	2 (5,1)	2 (7,4)	1 (7,1)
Amoxicilina-ácido clavulánico	3 (6,4)	0	3 (15,0)	1 (2,6)	1 (3,7)	0
Cefuroxima	3 (6,4)	2 (6,3)	1 (5,0)	1 (2,6)	0	0
Cefotaxima	3 (6,4)	2 (6,3)	1 (5,0)	1 (2,6)	0	0
Ceftriaxona	3 (6,4)	2 (6,3)	1 (5,0)	1 (2,6)	0	0
Gentamicina	0	2 (6,3)	0	4 (10,3)	0	0
Nitrofurantoina	1 (2,1)	0	2 (10,0)	0	0	0
Imipenem	0	0	0	0	1 (3,7)	0
Cefoxitina	0	0	1 (5,0)	0	0	0