

First record of the genus and species *Zefevazia peruana* (Boucomont, 1902) (Coleoptera: Bolboceratidae: Bolboceratinae) from Ecuador

Primer registro del género y la especie *Zefevazia peruana* (Boucomont, 1902) (Coleoptera: Bolboceratidae: Bolboceratinae) para Ecuador

Julián Clavijo-Bustos *¹

<https://orcid.org/0000-0002-7453-9792>
jclavijo@humboldt.org.co

Saúl S. Guillén²

<https://orcid.org/0000-0002-0749-8898>
saul.gonzalez-g@up.ac.pa

Manuel D. Barria³

<https://orcid.org/0000-0002-7045-0707>
manuel.barriag@up.ac.pa

*Corresponding author

1 Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Sección de entomología, Colecciones biológicas, Villa de Leyva, Boyacá, Colombia.

2 Universidad de Panamá, Facultad de Ciencias Naturales Exactas y Tecnología, Estudiante de la Escuela de Biología, Ciudad de Panamá, Panamá.

3 Universidad de Panamá (UP), Vicerrectoría de Investigación y Postgrado, Laboratorio de Estudios Biológicos de Artrópodos, Ciudad de Panamá, Panamá.

Citación

Clavijo-Bustos J, Guillén SS, Barria MD. 2022. First record of the genus and species *Zefevazia peruana* (Boucomont, 1902) (Coleoptera: Bolboceratidae: Bolboceratinae) for Ecuador. *Revista peruana de biología* 29(3): e22756 001- 004 (Agosto 2022). doi: <http://dx.doi.org/10.15381/rpb.v29i3.22756>

Presentado: 22/04/2022

Aceptado: 31/05/2022

Publicado online: 29/08/2022

Editor: Diana Silva

Abstract

The first report of the genus and species *Zefevazia peruana* (Boucomont, 1902) for Ecuador is presented. The species is previously known only from a few localities in northern Peru, and is now also recorded from Santa Elena, southern Ecuador. This report represents the northernmost record of the genus and species.

Resumen

Se presenta el primer reporte del género y la especie *Zefevazia peruana* (Boucomont, 1902) para Ecuador. La especie es conocida previamente de unas pocas localidades en el norte de Perú, y ahora también es registrada en Santa Elena, sur de Ecuador. Este reporte representa el registro más al norte del género y la especie.

Keywords:

Distribution; scarabs; Neotropics; taxonomy; Bolboceratinae.

Palabras clave:

Distribución; escarabajos; Neotrópico; taxonomía; Bolboceratinae.

Introduction

The subfamily Bolboceratinae belongs to the family Bolboceratidae, which was previously considered as a subfamily of Geotrupidae; however, Scholtz & Browne (1996) show the polyphyly between both taxa based on morphology and separated them as different families. Also, this fact has been partially demonstrated in recent molecular phylogenies of Coleoptera (*e.g.*, Ahrens et al. 2014, McKenna et al. 2015, Neita-Moreno et al. 2019). Despite the morphological and molecular evidence, there is no consensus, and in recent works this group of beetles is listed as Geotrupidae: Bolboceratinae (*e.g.*, Bouchard et al. 2011) or Bolboceratidae (*e.g.*, Scholtz & Grebennikov 2016).

In South America, Bolboceratinae (Coleoptera: Bolboceratidae) is constituted by eight genera and 42 species (Martínez 1976, Ide & Martínez 1993, 1994; Mondaca & Smith 2008, Carvalho & Vaz-de-Mello 2022). Four South American bolboceratine species belong to the genus *Zefevazia* Martínez, 1954, which is mainly distributed in Argentina and Paraguay (Martínez 1976, Mondaca & Smith 2008); however, *Zefevazia peruana* (Boucomont, 1902) was described only for Peru and so far, is the northernmost record of this genus. *Zefevazia* was originally described by Martínez (1954), later, the same author considered it as a subgenus of *Bolborhinum* Boucomont, 1911 (Martínez 1976). Finally, Mondaca & Smith (2008) revised *Bolborhinum* and elevated the status of its two subgenera: *Zefevazia* and *Pereirabolbus* Martínez, 1976.

We present the first record of *Zefevazia peruana* (Boucomont, 1902) for Ecuador extending the record of this genus further into northern South America.

Material and methods

The single examined specimen is deposited at the Section of the Entomology and Biological Collections, of the Instituto de Investigación de Recursos Biológicos Alexander von Humboldt - **IAvH-E**, at Villa de Leyva, Boyacá, Colombia.

Specimen photos were taken with a camera Canon EOS 5d Mark III with a macro lens Canon EF 100 mm f/2.8; genitalia images were taken with a camera Leica MC190HD attached to a stereomicroscope Leica S8 APO. Distribution map was elaborated with Tableau Public 2021, ArcGIS Pro and Adobe Illustrator 2020, and is based on specimen label information and records in literature.

Results

ZEFEVAZIA MARTÍNEZ, 1954

Zefevazia Martínez, 1954: 193. Type species: *Zefevazia ro-sascostai* Martínez, 1954 by original designation.

Diagnosis (*sensu* Mondaca & Smith 2008). Head in males with one frontal horn, females with one tubercle; antennal club semicircular. Pronotum in males with paired anteromedian excavations, which are marked basally by a transverse ridge; females with a small median concavity located behind the anterior margin of pronotal disc. Scutellum triangular. Elytron basally margined, with seven striae between medial suture and humeral swelling. Prosternal process reduced to absent. Metasternum pyriform, bulbous, with a long median carina; metasternal process wide, convex between mesocoxae. Mesocoxae distinctly separated. Male aedeagus subcylindrical, parameres short and subrectangular, truncated apically, apex wide.

Zefevazia peruana (Boucomont, 1902)

Bolboceras peruanus Boucomont, 1902: 185. Type locality: Grau, province de Thumbez, Pérou.

(Figs. 1–3)

Distribution. Peru (Boucomont 1902, Martínez 1976, Juárez-Noé & González-Coronado 2018), Ecuador (**new country record**) (Fig. 3).




Figure 1. *Zefevazia peruana* (Boucomont, 1902), male from Ecuador. (A) Dorsal view. (B) Ventral view. (C) Lateral view. (D) Frontal view. (E) Frontal dorsal view.

Material examined. ECUADOR, Santa Elena, La Libertad, Hotel Samarina, 02°13'S 80°52'W, 2 m, atraído por luz artificial, 01-31-xii.1997, Forero D. (1♂, IAvH-E-88458).

Records from literature. PERU: Piura [not specified which of the six localities sampled] (Juárez-Noé & González-Coronado 2018); Sullana, Hacienda Mallares (Martínez 1976); Tumbes, Grau (Boucomont 1902).

Diagnostic redescription . - Male (Figs. 1, 2). Length 13.6 mm. Color reddish-brown. *Head* with one cephalic vertical horn in the clypeal apex occupying the totality of the anterior edge; horn transversely compressed, with the apex truncated, slightly emarginated medially; anterior and posterior surface of cephalic horn coarsely punctate, near rugose, except at the very apex transversely smooth. Frons and vertex with sparsely small punctures. Labrum truncate, with lateral sides rounded, dorsal surface coarsely punctate, near rugose. *Pronotum* with two anteromedian excavations, space between excavations emarginate over the midline. Anteromedian

pronotal excavations delimited over the external edge in the anterior half, by a vertical carina with two lobules, the anterior longer than the posterior; anterior lobe of the carina curved to the inner side. Anterior angles of pronotum excavated, excavations located right after the anteromedian excavations and the vertical carina, dorsally covered with minute punctures. Dorsal surface of the posterior half and the midline in the anterior half of pronotum covered with round fine punctures separated by one to two their diameter; lateral punctures of the posterior half, near the vertical carinae, coarser and bigger, somewhat coalescent. Punctures in the surface of the anteromedian excavations slightly smaller and transverse, with a strigulose appearance. *Scutellum* triangular in shape, wider than long, surface with fine punctures. *Elytron* with 14 distinctly punctate striae, seven between median sutura and humeral swelling; punctures fine and separated by five times their diameter. Surface of interstriae sparsely covered with small punctures. *Anterior tibiae* with five teeth on the outer margin. *Aedeagus* and *genital capsule* as in Figure 2.




Figure 2. *Zefevazia peruana* (Boucomont, 1902), male terminalia. (A) Aedeagus, dorsal view. (B) Aedeagus, ventral view. (C) Genital capsule, dorsal view. (D) Genital capsule, ventral view.

Remarks. *Zefevazia peruana* was reported only from the Ecuadorian province of the Pacific dominion (*sensu* Morrone 2014, Morrone et al. 2022) (Fig. 3). The new record for Ecuador, also represents the northernmost record for the genus and species and the new record for the western Ecuador province, also in the Pacific dominion (Fig. 3). Furthermore, the species was reported in the Peruvian Department of Piura, in seasonally lowland dry forest (Juárez-Noé & González-Coronado 2018).

Despite that we were unable to study the female of the species, according to Martínez (1976), females of *Z. peruana* and its closely related species, *Zefevazia cantisani* (Martínez, 1976), could be distinguished by the shape of the clypeal horn (robust in the later), the lateral carina delimiting each anteromedian excavation (obsolete in the later), and surface of the anterolateral excavations with punctuations (more evident and closely distributed in the later).




Figure 3. Distribution of *Zefevazia peruana* (Boucomont, 1902) .

Discussion

Zefevazia peruana has been recorded in the departments of Piura and Tumbes in Peru, and in Santa Elena province in Ecuador. These territories correspond to the biogeographic Ecuadorian and Western Ecuador provinces, according to Morrone (2014) and Morrone et al. (2022). However, according with the biogeographical regionalisation of Udvardy (1975), correspond to the Ecuadorian Dry Forest province, which is the dominant vegetation type of the zone and the habitat in which the species has been collected (Juárez-Noé & González-Coronado 2018).

The other three species of *Zefevazia* are known from Argentina and Paraguay (Martínez 1976), therefore, *Z. peruana* is the northernmost species of the genus. Also, the new record from Ecuador now represents the northernmost record of the genus and species, which involve an extension of around 150 kilometers in straight line from the previous known closest record. Despite these data, the distribution of the species of *Zefevazia* are still poorly known and it is probable that important distribution extensions for all the species will occur in the future, as well as the possible discovery of new species.

Two genera of Bolboceratinae, each represented by one species, have been previously recorded from Ecuador: *Bolbapium baeri* (Boucomont, 1902) and *Parabolbapium aequatoriensis* Ide & Martínez, 1994 (Martínez 1976, Ide & Martínez 1994). The new record presented here raised to three the number of genera and species of the subfamily in the country. However, further detailed studies will show the composition and distribution of Bolboceratinae in Ecuador.

Literature cited

- Ahrens D, Schwarzer J, Vogler AP. 2014. The evolution of scarab beetles tracks the sequential rise of angiosperms and mammals. *Proceedings of the Royal Society B*, 281: 20141470. <http://dx.doi.org/10.1098/rspb.2014.1470>
- Bouchard P, Bousquet Y, Davies AE, Alonso-Zarazaga MA, Lawrence JF, Lyal CHC, Newton AF, Reid CAM, Schmidt M, Šlipiňský A, et al. 2011. Family-group names in Coleoptera (Insecta). *Zookeys* 88: 1-972. <https://doi.org/10.3897/zookeys.88.807>
- Boucomont JA. 1902. Description sommaire de quelques Bolbocerina (Col.). *Bulletin de la Société entomologique de France*: 184-185.
- Carvalho EC, Vaz-de-Mello FZ. 2022. A taxonomic revision of *Bolbapium* Boucomont, 1910 (Coleoptera: Scarabaeoidea: Geotrupidae). *Journal of Natural History* 56(13): 769-828. <https://doi.org/10.1080/00222933.2022.2092429>
- Juárez-Noé G, González-Coronado U. 2018. Lista de Coleópteros (Insecta: Coleoptera) de la región Piura, Perú. *Folia Entomológica Mexicana (nueva serie)* 4(1): 1-27. <https://doi.org/10.3989/graellsia.2019.v75.233>
- Ide S, Martínez A. 1993. Two new species of *Bolbapium* Boucomont, 1910 (Coleoptera, Geotrupidae, Bolboceratinae). *Revista Brasileira de Entomologia* 37(2): 369-385.
- Ide S, Martínez A. 1994. Description of *Parabolbapium aequatoriensis*, new genus and new species from Ecuador (Coleoptera, Geotrupidae). *Revista Brasileira de Entomologia* 38(2): 435-446.
- Martínez A. 1954. Un nuevo género de Bolboceratini Neotropical (Coleoptera; Scarabaeidae). *Revista Brasileira de Entomologia* 12: 193-204.
- Martínez A. 1976. Contribución al conocimiento de los Bolboceratini sudamericanos (Coleoptera, Scarabaeidae, Geotrupinae, Bolboceratini). *Estudia Entomológica* 19(1-4): 531-551.
- McKenna DD, Wild AL, Kanda K, Bellamy CL, Beutel RG, Caterino MS, Farnum CW, Hawks DC, Ivie MA, Jameson ML, et al. 2015. The beetle tree of life reveals that Coleoptera survived end-Permian mass extinction to diversify during the Cretaceous terrestrial revolution. *Systematic Entomology* 40(4): 835-880. <https://doi.org/10.1111/syen.12132>
- Mondaca J, Smith ABT. 2008. A revision of the southern South American genus *Bolborhinum* Boucomont (Coleoptera: Geotrupidae: Bolboceratinae). *Zootaxa* 1794: 1-48. <https://doi.org/10.11646/ZOOTAXA.1794.1.1>
- Morrone JJ. 2014. Biogeographic regionalisation of the Neotropical region. *Zootaxa* 3782: 001-110. <https://doi.org/10.11646/zootaxa.3782.1.1>
- Morrone JJ, Escalante T, Rodríguez-Tapia G, Carmona A, Arania M & Mercado-Gómez JD. 2022. Biogeographic regionalization of the Neotropical region: new map and shapefile. *Anais da Academia Brasileira de Ciências* 94(1): e20211167. <https://doi.org/10.1590/0001-376520220211167>
- Neita-Moreno JC, Agrain F, Eberle J, Ahrens D & Pereyra V. 2019. On the phylogenetic position and systematics of extant and fossil Aclopininae (Coleoptera: Scarabaeidae). *Systematic Entomology* 44(4): 709-727. <https://doi.org/10.1111/syen.12366>
- Scholtz CH, Browne DJ. 1996. Polyphyly in the Geotrupidae (Coleoptera: Scarabaeoidea): a case for a new family. *Journal of Natural History* 30: 597-614. <http://dx.doi.org/10.1080/00222939600770311>
- Scholtz CH, Grebennikov VV. 2016. Scarabaeoidea Latreille, 1802. In: Beutel RG, Leschen RAB, eds. *Coleoptera, Beetles. Morphology and Systematics*. Coleoptera, beetles. Morphology and systematics. Walter de Gruyter GmbH & Co KG. Pp. 443-525.
- Udvardy MDF. 1975. A classification of the biogeographical provinces of the world. IUCN Occasional Paper No 18. International Union for Conservation of Nature and Natural resources, Switzerland. 49 pp.

Agradecimientos / Acknowledgments:

This publication was partially supported by the Instituto de Investigación de Recursos Biológicos Alexander von Humboldt. We thank J. C. Neita Moreno, curator of the IAvH-E, for allowing the study of the specimen from Ecuador.

Conflicto de intereses / Competing interests:

The authors declare no conflict of interest.

Rol de los autores / Authors Roles:

JCB: Conceptualización, Investigación, Escritura-Preparación del borrador original, Redacción-revisión y edición
SSG: Conceptualización, Investigación, Redacción-revisión y edición.
MDB: Conceptualización, Redacción-revisión y edición.

Fuentes de financiamiento / Funding:

This research received no specific funding from any agency, commercial or not-for-profit sectors.

Aspectos éticos / legales; Ethics / legal:

Authors declare that they did not violate or omit ethical or legal norms in this research.