

DETERMINATION OF ANTHOCYANINS, TOTAL PHENOLS AND ANTIOXIDANT CAPACITY OF *RUBUS ROBUSTUS C. PRESL*

Daniela Camacho-Valencia^{a,b}, Stamber Alvaro Ramírez- Revilla ^{*b},
José Antonio Villanueva-Salas^a

ABSTRACT

Natural antioxidants, such as fruits and vegetables, can neutralize an excess of free radicals in the organism and prevent certain degenerative diseases. Blackberries are known to exhibit antioxidant properties, but this fruit is seldom consumed or commercialized in Peru. The objective of this research study was to determine the total anthocyanins (TA), total phenolic compounds (TPC) and antioxidant capacity of lyophilized *Rubus robustus C. Presl* (wild blackberries) in the region of Arequipa - Peru. The pH-differential method, the Folin-Ciocalteu method and the free radical 2,2-diphenyl-1-picrylhydrazyl hydrate (DPPH) method were used to this end. Results showed a TA content of 52 ± 1.86 mg cyanidin 3-glucoside/100 g of lyophilized *Rubus robustus C. Presl*. Furthermore, the TP content was of 219.11 ± 8.23 mg gallic acid/100 g of the lyophilized fruit. The percentage of DPPH discoloration of two samples (1.0 and 2.0 mg/mL) of the lyophilized fruit was determined, obtaining a percentage of 35.89 ± 4.43 and $67.23 \pm 3.30\%$, respectively. The IC₅₀ value found was 1.5 mg/mL. Wild blackberries in Arequipa, Peru were therefore shown to possess an antioxidant capacity, so a diet rich in this type of fruit should be promoted as a nutraceutical.

Keywords: blackberries; anthocyanins; total phenols; DPPH; antioxidant.

DETERMINACIÓN DE ANTHOCIANINAS, FENOLES TOTALES Y CAPACIDAD ANTIOXIDANTE DE *RUBUS ROBUSTUS C. PRESL*

RESUMEN

Los antioxidantes naturales, como las frutas y vegetales, pueden neutralizar un exceso de radicales libres en el organismo y prevenir ciertas enfermedades degenerativas. Se sabe que las zarzamoras exhiben propiedades antioxidantes, pero esta fruta es rara vez consumida o comercializada en Perú. El objetivo de este estudio de investigación fue determinar el total de antocianinas (TA), fenoles totales (TP) y capacidad antioxidante del liofilizado de *Rubus robustus C. Presl* (zarzamoras silvestres) en la región de Arequipa - Perú. Para ello, se utilizaron el método diferencial de pH, el método Folin-Ciocalteu y el método de radicales

^a Universidad Católica de Santa María, Urb. San José s/n Umacollo, Arequipa 04001, Perú

^b Universidad Tecnológica del Perú, Tacna y Arica 160, Arequipa 04001, Perú

*corresponding author: sramirezr@utp.edu.pe

libres de 2,2-difenil-1-picrilhidracilo (DPPH). Los resultados mostraron un contenido de TA de 52 ± 1.86 mg de cianidina 3-glucósido/100 g de *Rubus robustus* C. Presl liofilizado. Además, el contenido de TP fue de 219.11 ± 8.23 mg de ácido gálico/100 g de la fruta liofilizada. Se determinó el porcentaje de decoloración DPPH de dos muestras (1.0 y 2.0 mg/mL) de la fruta liofilizada, obteniendo un porcentaje de 35.89 ± 4.43 y $67.23 \pm 3.30\%$, respectivamente. El valor de IC50 encontrado fue de 1.5 mg/mL. Por lo tanto, se demostró que las zarzamoras silvestres en Arequipa, Perú, poseen una capacidad antioxidante, por lo que una dieta rica en este tipo de frutas debe promoverse como nutracéutico.

Palabras clave: zarzamoras; antocianinas; fenoles totales; DPPH; antioxidante.

INTRODUCTION

Free radicals are a molecular species with an unpaired electron in an atomic orbital, causing them to be highly reactive and capable of altering different molecules in the body, resulting in the progression of a number of diseases and also in aging¹. There are many sources of free radicals, including enzymes such as xanthine oxidase, cyclooxygenases, lipoxygenases, myeloperoxidases, cytochrome P450 monooxygenase, uncoupled nitric oxide synthase (NOS), peroxidases and NADPH oxidase².

Antioxidants can prevent cellular damage resulting from chemical reactions of free radicals by neutralizing and stabilizing them³. Different studies have reported the effects of natural antioxidants in diseases related to stress oxidation, such as cancer or atherosclerosis^{4,5}.

Blackberries belong to the genus *Rubus* of the rose family (Rosaceae), it is a berry fruit which consists of numerous black juicy drupelets, each with a single seed around a central stem. It is commonly found in North America and Europe. Blackberries are eaten fresh or in other food products such as ice cream, jams and pies⁶. Blackberries are known to be rich in anthocyanins and other phytochemicals, such as flavonols, phenolic acids, ellagic acid, vitamins C and E, folic acid and b-sitosterol⁷. The phenolic composition, antioxidant capacity and anthocyanin content of different species of blackberries⁸⁻¹³ have been studied using different determination techniques most of all concluding that this fruit has an antioxidant capacity due to their high anthocyanin content, more precisely from cyanidin-3-glucoside¹⁴. In Peru, blackberries are poorly grown and often imported but this fruit can grow by itself like tare with a source of water nearby.

We emphasize the importance of publishing information about blackberries as antioxidants in Peru since there are few reports available on the antioxidant capacity of this fruit in this region. This research will be helpful for greater understanding the value of blackberries to the health of Peruvian consumers and for encouraging its consumption. The aim of this research study, therefore, was to determine total anthocyanins (TA), total phenols (TP) and antioxidant capacity of *Rubus robustus* C. Presl (wild blackberries) in Arequipa, Peru.

EXPERIMENTAL

This research was carried out in the Proyecto Mercurio Laboratory (H-202) at the Universidad Católica de Santa María and in the Calidad de Aguas Laboratory at the Universidad Tecnológica del Perú in Arequipa.

Fruit Sample

Mature wild blackberries were collected from adult wild plants in the Chilina Valley in the spring (between September and October) as shown in Figure 1. Gathering was carried out by direct hand harvesting of the fully ripe fruits and they were selected by uniformity of size and color and stored in polyethylene terephthalate (PET) containers. Fruit samples were sent to the Proyecto Mercurio Laboratory, where they were carefully washed with deionized water and kept refrigerated at 4 ± 1 °C. Later, the samples were blended and filtrated to eliminate any solid material before being freeze-stored. Afterwards, the samples were lyophilized in a FreeZone 2.5 Liter Benchtop Freeze Dry System at the Universidad Nacional San Agustín of Arequipa. Finally, the lyophilized samples were placed in PET containers and stored in an environment free of humidity and protected from direct light. A sample of the adult plant and fruit were sent to the Herbarium Arequipense at the Universidad Nacional San Agustín of Arequipa, which identified it as *Rubus robustus* C. Presl.



Figure 1. *Rubus robustus* C. Presl collected from the Chilina Valley

Sample preparation

The lyophilized sample (100 mg) was treated with 1 mL of high-performance liquid chromatography (HPLC) grade methanol and sonicated in a Branson 2510-E sonicator for 10 min. The homogenate was then centrifuged in a Centra CL 2 Thermo IEC centrifuge at 4000 rpm for 15 min at room temperature. The supernatant was separated and the remaining insoluble fraction was washed again with 1 mL of HPLC grade methanol. This process was repeated three times and the supernatants were combined. Finally, a final volume of 10 mL was obtained with HPLC grade methanol and maintained at 4 ± 1 °C.

Reagents

Follin-Ciocateau reagent P.A (Merck), gallic acid P.A (Sigma Aldrich), sodium carbonate P.A (Merck), potassium chloride P.A (Merck), sodium acetate P.A (Merck), 6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox) standard (CALBIOCHEM), DPPH P.A

(CALBIOCHEM), HPLC grade methanol (MERCK), deionized water (18.2 mΩ) obtained from a Barnstead Easypure II water purification system.

Determination of total anthocyanins

The pH differential method has been used by many authors to determine the anthocyanins of berries¹¹⁻¹³ where a change in pH, anthocyanins go through a reversible structural transformation demonstrated by different absorbance spectra. At pH 1.0, the red oxonium form appears and at pH 4.5 the colorless hemiketal form predominates¹⁵. Absorbance was measured with a CARY60 UV –VIS spectrophotometer (Agilent Technologies) at 510 and 700 nm at different pHs to calculate the final absorbance using Eq. 1

$$A = (A_{510} - A_{700})_{pH\ 1} - (A_{510} - A_{700})_{pH\ 4.5} \quad (1)$$

The TA was calculated according to Eq. 2:

$$(TA) = \frac{A \times MW \times DF \times 100}{\epsilon \times l} \quad (2)$$

Where: A= Absorbance, MW= molecular weight, DF= dilution factor, ε= extinction coefficient. Data were calculated with the MW (449.20) and the extinction coefficient for cyanidin-3-glucoside (29 600) and expressed as mg cyanidin/100 g lyophilized *Rubus robustus* C. Presl.

Determination of total phenolics compounds

The TP was determined by the Folin-Ciocalteu colorimetric method, first described by Singleton and Rossi (1965)¹⁶ based on the production of a blue phosphotungstic-phosphomolydic complex. This method is widely used to estimate the total phenolic content in many samples such as fruits, beverages, herbs and other plant extracts¹⁷⁻¹⁹.

A calibration curve was prepared with gallic acid ranging from 10 to 60 ppm. Different concentrations of gallic acid were mixed with 0.25 mL of Folin-Ciocalteu reagent in a 10 mL volumetric flask at room temperature. Then, 2 mL of a 20% sodium carbonate solution was added and finally deionized water was added to complete 10 mL. This solution was mixed in an Analog mixer vortex and maintained for 2 h in dark conditions. Finally, total sample phenols were evaluated with a CARY60 UV –VIS spectrophotometer (Agilent Technologies) at 750 nm.

For the determination of TP of the fruit sample, the same procedure was done using 0.2 mL of the sample previously prepared. TP was expressed as mg of gallic acid equivalent (GAE)/100 g lyophilized *Rubus robustus* C. Presl.

Antioxidant assay: Determination of the DPPH free radical scavenging activity

The DPPH free radical method is based on the measurement of the scavenging capacity of antioxidants towards a free radical²⁰ and it is used in various studies to determine the antioxidant capacity of plants²¹⁻²³. When a DPPH solution is mixed with a substance such as an antioxidant that can donate a hydrogen atom, it gives rise to a reduced form with the loss of its violet color²⁴.

A stock solution of Trolox at a concentration of 1 mmol/L was prepared with HPLC grade methanol in a 10 mL volumetric flask and stored in the fridge until use. A working solution of DPPH at a concentration of 2.5 mmol/L was prepared with HPLC grade methanol and was kept in dark conditions to prevent its deterioration. This solution was prepared on each day of the analysis.

A calibration curve was prepared using different volumes of the stock solution of Trolox to obtain a concentration range of 0.02–0.10 mmol/L, 100 uL of the DPPH working solution and HPLC grade methanol to achieve a volume of 1 mL. To prepare the negative control, DPPH was mixed with methanol instead of Trolox. Finally, for the analysis of the sample, two different volumes (100 and 200 uL) of the fruit samples prepared earlier were taken, 100 uL of the DPPH and HPLC grade methanol were added to obtain a volume of 1 mL and filtrated in an Anotop filter before injecting them to the HPLC. The absorbance was measured at 517 nm every 20 min. The samples and standards were analyzed. The peaks identification was performed by the comparison of the retention times with those of referential standards.

The analysis was performed on a LaChrom-Hitachi HPLC, consisting of an L-7100 pump, a 20 uL manual injector (loop) and a L-7400 UV detector. An EZ Chrom Elite data station and a personal computer were used as software sources for data storage and evaluation. The analytical column used was a Chromolith RP-18e of 100–4.6 mm, with internal particle diameter of 2 um. A 5–4.6 mm Chromolith RP-18e pre-column was used to protect the analytical column. Chromatographic separation was performed using, as a mobile phase methanol: water (80:20) with a total flow rate at 1.0 mL/ min. An absorbance detector was used at 517 nm.

The results were expressed as a percentage decrease of color intensity (% inhibition of the free radical) related to the negative control sample such as Eq. 3:

$$(\% \text{ inhibition}) = \frac{A_{\text{negative control}} - A_{\text{sample}} * 100}{A_{\text{negative control}}} \quad (3)$$

where: A_{negative} = area of the negative control and A_{sample} = area of the sample.

Determination of the IC₅₀

The IC₅₀ is the concentration of an antioxidant that inhibits by 50% the DPPH free radical activity. The IC₅₀ is inversely proportional to the antioxidant activity where a lower value indicates a higher effectiveness of the antioxidant²³. To determine the IC₅₀, a calibration curve was prepared using serial dilutions made from the stock solution of the fruit sample to obtain solutions with concentrations of 3000, 2000, 1500, 1000, 800, 500 and 200 µg/mL, mixing them with 100 uL of the DPPH working solution and HPLC grade methanol. The working solution of DPPH 2.5 mmol/L was prepared as previously explained. To prepare the negative control, DPPH was just mixed with methanol. Finally, the absorbance was measured at 517 nm on a GENESYS UV –VIS spectrophotometer (Thermoscientific)

Statistical analysis

The data obtained was analyzed on the basis of a random design using the general linear model with five replicates and by analysis of variance (ANOVA). All statistical tests were performed with a level of significance of 0.05. Significant differences among means were determined by Tukey's test at 5% level of significance.

RESULTS AND DISCUSSION

Determination of TA and TPC

Total anthocyanins and total phenolic contents are presented in Table 1. Lyophilized *Rubus robustus* C. Presl presented 52 ± 1.86 mg of total anthocyanins whereas Croge et al. (2019)¹², presented a value between 80 a 134 mg /100 g of fresh fruit for different blackberries species hence the fruit extract possesses a higher anthocyanin content possibly due to the extraction method. What is more, in a nearby country such as Brazil, the content of phenolic compounds in blackberry was reported with values ranging 234 to 800 mg GAE/ 100 mg of fresh fruit¹²⁻²⁵, where as the lyophilized sample of *Rubus robustus* C. Presl presented 219.11 ± 8.23 mg GAE/100 mg indicating the climate and the temperature influences in the content of these compounds. In addition, in Peru, values of TPC for *Rubus fruticosus* L 400.67 mg GAE /100 g of fresh fruit were found²⁶ showing a difference with the lyophilized *Rubus robustus* C. Presl which could be due to the species of Rubus, the type of sample and the growing conditions.

Table 1. Contents of total anthocyanins and total phenolic compounds of lyophilized *Rubus robustus* C. Presl

Total anthocyanins mg/100 g lyophilized
<i>Rubus robustus</i> C. Presl
52.00 ± 1.86
Total phenolic compounds mg/100 g
lyophilized <i>Rubus robustus</i> C. Presl
219.11 ± 8.23

Antioxidant assay: Determination of the DPPH free radical scavenging activity


Eq. 3 was used to calculate the percentage of inhibition of DPPH. Table 2 presents the results of the percentage of inhibition of DPPH in 5 different repetitions using different concentrations of Trolox. As seen in Table 2, if the concentration of the antioxidant increases, so does the percentage of inhibition of the free radical.

Table 2. Percentage of inhibition of DPPH using different concentrations of Trolox.

Concentrations of Trolox (mmol/mL)	Percentage of inhibition (%)
0.02	13.67 ± 3.34
0.04	28.35 ± 5.43
0.06	42.81 ± 7.06
0.08	56.4 ± 6.04
0.1	71.96 ± 7.67

Results show that 1.0 and 2.0 mg/mL of the lyophilized sample present a percentage of inhibition of 35.89 ± 4.43 and $67.23 \pm 3.30\%$, respectively. In another study, the DPPH discoloration percentages of different fresh berries was determined, where wild blackberries had a percentage of 87.5%²⁷, showing that fresh blackberries have more antioxidant capacity than the lyophilized samples in this study with 67%. Furthermore, in a study conducted with *Rubus laciniatus cv. Hull* (thornless blackberry) in China, at 2 mg/mL, blackberry extracts could scavenge nearly all DPPH radicals (95.37%)²⁸ whereas in Peru, with *Rubus robustus C. Presl* at the same concentration, the percentage of inhibition was 67%, demonstrating a difference in the species and in the location. What is more, the scavenging capacity of five blackberry cultivars produced in Brazil²⁹ was determined, ranging from 53 to 74%, proving that *Rubus robustus C. Presl* has an average antioxidant capacity.

Figure 2 illustrates the chromatograms of three samples: A) DPPH + methanol, B) DPPH + methanol + Trolox (0.06 mmol/mL) and C) DPPH + methanol + sample (1.0 mg/mL). As seen in these chromatograms, the retention time is approximately 2.91 min and the total runtime was 5 min. When comparing chromatogram A and B, chromatogram B shows that a Trolox concentration (in this case 0.06 mmol/mL) decreases the area of the DPPH peak, mainly because the Trolox solution is used as an antioxidant. In chromatogram C, the sample is proven to contain antioxidant capacity as the peak area of DPPH decreases compared to chromatogram A, with an effect similar to that of the Trolox solution used.

**Figura 2.** Chromatograms: A) DPPH + methanol, B) DPPH + methanol + Trolox (0.06 mmol/mL) and C) DPPH + methanol + sample (1.0 mg/mL blackberry sample)

Determination of the IC50

Table 3 summarizes the DPPH radical scavenging activity of lyophilized *Rubus robustus* C. Presl. All experiments were conducted in triplicate (n=3) and reported as the mean of three values together with standard deviation (\pm SD). The IC50 value for lyophilized *Rubus robustus* C. Presl was found to be approximately 1.5 mg/mL hence proving that this species of blackberry in Peru possess an antioxidant activity.

Tabla 3. summarizes the DPPH radical scavenging activity of lyophilized *Rubus robustus* C. Presl

Concentrations (μg/mL)	Percentage of inhibition (%)
200	10.39 \pm 3.15
500	19.18 \pm 1.10
800	25.24 \pm 0.95
1000	38.15 \pm 4.64
1500	53.52 \pm 2.60
2000	57.06 \pm 3.56
3000	58.42 \pm 2.10

Other blackberries species like the Brazilian Blackberry Xavante and Blackberry Cherokee have demonstrated a high IC5025. In addition, when comparing different small berries in Italy, blackberries have evidenced to be one of the best natural antioxidants¹³.

CONCLUSIONS

Wild blackberries collected from the Chilina Valley in Arequipa, Peru, exhibited antioxidant capacity demonstrated by their content of anthocyanins (52 ± 1.86 mg cyanidin 3-glucoside/100 g of lyophilized *Rubus robustus* C. Presl) and phenols (219.11 ± 8.23 mg gallic acid/100 g of the lyophilized fruit) as well as by the antioxidant assay with a percentage of 35.89 ± 4.43 and $67.23 \pm 3.30\%$, respectively. In addition, the IC50 value found was 1.5 mg/mL. A diet rich in this fruit could provide a good source of antioxidants, and therefore it should be encouraged in Peru. Blackberries are not currently extensively grown in Arequipa, but their consumption may have potential for use in the development of nutraceuticals for regional and international markets. Furthermore, they could be used as functional food ingredients or supplements as a benefit to human health. Further studies should be done to compare the antioxidant capacity in cultivated and wild blackberries in the same area and in other areas of Arequipa, Peru.

REFERENCIAS BIBLIOGRÁFICAS

1. Lobo V, Patil A, Phatak A, Chandra N. Free Radicals, Antioxidants and Functional Foods: Impact on Human Health. *Pharmacogn Rev.* 2010; 4(8): 118–126.
2. Siti HN, Kamisah Y, Kamsiah J. The role of oxidative stress, antioxidants and vascular inflammation in cardiovascular disease (a review). *Vascul Pharmacol.* 2015; 71: 40–56.
3. Young IS, Woodside JV. Antioxidants in health and disease. *J Clin Pathol.* 2001; 54(3): 176–186.
4. Seeram NP. Berry Fruits for Cancer Prevention: Current Status and Future Prospects. *J Agric Food Chem.* 2008, 56(3): 630–635.
5. Seeram NP, Adams LS, Zhang Y, Lee R, Sand D, Scheuller HS, et al. Blackberry, Black Raspberry, Blueberry, Cranberry, Red Raspberry, and Strawberry Extracts Inhibit Growth and Stimulate Apoptosis of Human Cancer Cells In Vitro. *J Agric Food Chem.* 2006; 54(25): 9329–9339.
6. Padmanabhan P, Correa-Betanzo J, Paliyath G. Berries and Related Fruits. *Encyclopedia of Food and Health.* Oxford: Academic Press. 2018; 364-371. doi:10.1016/B978-0-12-384947-2.00060-X.
7. Bowen-Forbes CS, Zhang Y, Nair MG. Anthocyanin content, antioxidant, anti-inflammatory and anticancer properties of blackberry and raspberry fruits. *J Food Compos Anal.* 2010; 23(6): 554–560.
8. Lee SG, Vance TM, Nam TG, Kim DO, Koo SI, Chun OK. Contribution of Anthocyanin Composition to Total Antioxidant Capacity of Berries. *Plant Foods Hum Nutr.* 2015; 70(4): 427–432.
9. Kostecka-Gugała A, Ledwożyw-Smołeń I, Augustynowicz J, Wyzgolik G, Kruczek M, Kaszycki P. Antioxidant properties of fruits of raspberry and blackberry grown in central Europe. *Open Chem.* 2015; 13(1): 1313–1325.
10. Van de Velde F, Grace MH, Esposito D, Pirovani MÉ, Lila MA. Quantitative comparison of phytochemical profile, antioxidant, and anti-inflammatory properties of blackberry fruits adapted to Argentina. *J Food Compos Anal.* 2016; 47: 82–91.
11. Prior RL, Sintara M, Chang T. Multi-radical (ORACMR5) antioxidant capacity of selected berries and effects of food processing. *J Berry Res.* 2016; 6(2): 159-173.
12. Croge C, Cuquel F, Matumoto-Pintro P, Biasi L, Bona C. Antioxidant Capacity and Polyphenolic Compounds of Blackberries Produced in Different Climates. *HortScience.* 2019; 54: 2209-2213.
13. Zorzi M, Gai F, Medana C, Aigotti R, Morello S, Peiretti P.G. Bioactive Compounds and Antioxidant Capacity of Small Berries. *Foods.* 2020; 9(5): 623. doi: 10.3390/foods9050623.
14. Fan-Chiang HJ, Wrolstad RE. Anthocyanin Pigment Composition of Blackberries. *J Food Sci.* 2005; 70(3): C198–C202.
15. Giusti MM, Wrolstad RE. Characterization and Measurement of Anthocyanins by UV-Visible Spectroscopy. *Curr Protoc Food Anal Chem.* 2001; 00(1): F1.2.1–F1.2.13. doi: 10.1002/0471142913.faf0102s00
16. Singleton V Rossi, J. Colorimetry of Total Phenolic Compounds with Phosphomolybdic-Phosphotungstic Acid Reagents. *Am J Enol Vitic.* 1965; 16(3): 144-158.

17. Margraf T, Karnopp AR, Rosso ND, Granato D. Comparison between Folin-Ciocalteu and Prussian Blue Assays to Estimate The Total Phenolic Content of Juices and Teas Using 96-Well Microplates. *J Food Sci.* 2015; 80(11): C2397–C2403. doi:10.1111/1750-3841.13077
18. Abramovič H, Grobin B, Poklar Ulrih N, Cigić B. Relevance and Standardization of In Vitro Antioxidant Assays: ABTS, DPPH, and Folin–Ciocalteu. *J Chem.* 2018; 2018: 4608405. doi: 10.1155/2018/4608405
19. Benabderrahim MA, Yahia Y, Bettaieb I, Elfalleh W, Nagaz K. Antioxidant activity and phenolic profile of a collection of medicinal plants from Tunisian arid and Saharan regions. *Ind Crops Prod.* 2019; 138: 111427. doi:10.1016/j.indcrop.2019.05.076.
20. Brand-Williams, W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *Food Sci Technol.* 1995; 28(1): 25–30.
21. Galvez MAC. Evaluation of DPPH Free Radical Scavenging Activity and Phytochemical Screening of Selected Folkloric Medicinal Plants in Tinoc, Ifugao, Cordillera Administrative Region, Philippines. *Int J Sci Res Publ.* 2015; 5(12): 440-445.
22. Patel R, Patel Y, Kunjadia P, AnjuKunjadia. DPPH free radical scavenging activity of phenolics and flavonoids in some medicinal plants of India. *Int J Curr Microbiol Appl Sci.* 2015; 4(1): 773-780.
23. Manoharan KP, Lehlohonolo IS, Sibusisiwe M. DPPH radical scavenging activity of extracts from *Rhamnus prinoides*. *J Med Plant Res.* 2019; 13(15): 329–334.
24. Kedare SB & Singh RP. Genesis and development of DPPH method of antioxidant assay. *J Food Sci Technol.* 2011; 48(4): 412–422.
25. Denardin CC, Hirsch GE, da Rocha RF, Vizzotto M, Henriques AT, Moreira JCF, et al. Antioxidant capacity and bioactive compounds of four Brazilian native fruits. *J Food Drug Anal.* 2015; 23(3) : 387–398.
26. Valencia Sullca CE, Guevara Pérez A. Variación de la capacidad antioxidante y compuestos bioactivos durante el procesamiento del néctar de zarzamora (*Rubus fruticosus* L.). *Rev Soc Quím Perú.* 2013; 79(2), 116-125.
27. Guerrero CJ, Ciampi PL, Castilla CA, Medel SF, Schalchli SH, Hormazabal UE, Bensch TE, Alberdi LM. Antioxidant Capacity, Anthocyanins, and Total Phenols of Wild and Cultivated Berries in Chile. *Chilean J Agric Res.* 2010; 70(4): 537–544.
28. Huang WY, Zhang HC, Liu WX, Li CY. Survey of antioxidant capacity and phenolic composition of blueberry, blackberry, and strawberry in Nanjing. *J Zhejiang Univ Sci B.* 2012; 13(2): 94–102.
29. Celant VM, Braga GC, Vorpapel JA, Salibe AB. Phenolic Composition and Antioxidant Capacity of Aqueous and Ethanolic Extracts of Blackberries. *Rev Bras Frutic.* 2016; 38(2): e-411. doi:10.1590/0100-29452016411.